

Boron Applications for Increased Apple Production

- Boron is essential for all plant growth. It aids in the transfer of sugars and nutrients from leaves to fruit, and increases pollination and seed development.
- Apples require an adequate supply of available boron, especially during flower formation and fruit set.
- Foliar sprays of *Solubor*® or *Solubor*® DF during the dormant season, pre-pink and early bud stages will ensure an adequate supply of boron during the critical stages of flowering and fruit development.
- Apples are sensitive to over-applications of boron, and boron toxicity can result if the recommended boron application methods and rates are not carefully followed.

Apples appear to grow best in sandy soils, but these soils also tend to be low in available boron (B), which is a key nutrient for tree growth – especially fruit set and quality.

While B is essential for all stages of plant growth, an available supply is most important during flowering and fruit development. This is because fruit trees have difficulty in transporting enough B to new flower buds, so foliar spray applications of *Solubor* at the pre-pink/pink stage will ensure an adequate supply of B during this critical reproductive stage. Cell wall strength, cell division, fruit and seed development and sugar transport are some of the plant functions related to B.

Available B in the soil is mainly associated with organic matter content, which must be mineralized to release B for plant uptake. While B requirements for optimum plant nutrition are low compared with those of the primary nutrients, the need for B is especially significant in fruit development.

Deficiency Symptoms

Because B is vital to flower formation and fruit set, a decrease in B supply during this critical stage can result in decreased yields. Boron deficiency results in reduced tree vigor and growth, and a severe deficiency may cause twig die-back, the formation of leaf rosettes, and bud drop.

Fruit “corking” (darkening of internal tissues) is another B-deficiency symptom. “Bitter pit” or tissue breakdown is another symptom because B is involved in the movement of calcium in plant tissues.

Soil Tests and Plant Analyses

Boron deficiencies may be suspected on light-textured soils where organic matter content is low, on soils with a pH above 6.0, and on recently limed soils. Soil testing and plant analyses are both helpful in assessing the potential B-supplying capacity of the soil and the current B status of the growing plant.

The critical level of hot-water-soluble B for apples in most soils is less than 0.5 ppm, depending on the soil pH, organic matter content and texture. The critical level of B in the top mature apple leaves is about 25 ppm, and the ideal range is 35-50 ppm.

Trees with leaf B contents below the critical level should be sprayed one or more times with *Solubor* before flower formation and fruit set, or with dormant sprays when flower buds begin development for the next crop. Trees with leaf B levels higher than 50 ppm should not be sprayed.

Recommendations for Apples

Yield responses to applied B are inconsistent and seasonal, probably due to environmental effects on growth. However, both yield and quality of apples may be improved with B fertilization because available B levels are low in some soils.

Apples are sensitive to over-applications of B, and toxicity can result if the recommended B application rates are not carefully followed.

Boron should be applied for apples, especially on sandy soils in high rainfall regions or with over-irrigation, because soluble B can be easily leached from the root zone.

Response to applied B generally is greatest when there are adequate supplies of other nutrients. While B recommendations for apples vary from state to state, two general types of recommendations are commonly given.

Boron Recommendations for Apples	
Marginal soil test B and/or leaf analyses, or dry weather during critical stages:	Low soil test B and a prior history of B response:
Foliar sprays at rates of 2.5 - 5.0 lbs. of <i>Solubor</i> /acre (0.5 - 1.0 lbs. of B/acre) at tight cluster to pink/white bud stage and also 7 - 10 days after petal fall. <i>Solubor</i> can be applied alone or with insecticides to plants. Post-harvest sprays, applied when leaves are still green, also can give a nutrient boost to buds developing for the next crop.	An early season soil application of 15 - 20 lbs. of <i>Granubor</i> /acre (2 - 3 lbs. of B/acre) surface broadcast, plus foliar sprays at 2.5 - 5.0 lbs. of <i>Solubor</i> /acre (0.5 - 1.0 lbs. of B/acre) per spray applied at a tight cluster to pink/white bud stage, and also at 7 - 10 and 25 - 30 days after petal fall.

For more information:

Call US Borax at
1 (800) 699 9005

Access our Fax on Demand
System at 1 (800) 472 9768

Visit our website at
www.borax.com/agriculture

