

B

Boron in forestry production

Boron deficiency is the most common micronutrient (4) limitation in forest plantations. It occurs in many countries, particularly in exotic plantations of eucalypts and pines, but also in plantations and natural stands of native species on soils altered by macronutrient fertilization, liming, fire or erosion.

Deficiency symptoms

Deficiency symptoms are usually characteristic but can be confounded by variable foliar concentrations, erratic occurrence, and possible climatic damages. Boron deficiency symptoms (4) on pines vary by species and may be distinctly seasonal in occurrence and are often environmental stress induced. Spring growth from unaffected buds appears normal, and it is not until midsummer or later that leading shoots show clearly recognizable symptoms. Limited supplies or even brief interruptions (such as those caused by drought) in absorption of boron can cause irreversible damage to rapidly growing shoots. A split leader is often the outcome.

Pinus sylvestris – (a uninodal species) exhibits abundant resin flow and various disturbances in apical dominance as the first external evidences of boron deficiency. The terminal bud may be small, malformed, retarded in expansion or dead. Adjacent lateral buds may or may not be similarly affected. Swelling, cracking, bending or dying of the leader occurs as in other pine species and so also darkening and formation of cavities in the pith. Needles near the affected tips are often short and deformed, either dark green, or discolored.

Other symptoms noted in various species are:

Pinus strobus and *Pinus sylvestris* – Dieback of terminal growing points. Tips of the primary needles turn light yellow/orange in color and have light brown edges.

Pinus sylvestris – Seedling are short with thick, succulent and somewhat fragile roots. The buds, which are small, remain moribund and the young needles chlorotic. Malformation of the needles is common.

Pinus radiata – Shoot and tip dieback, dead tops and malformation, brown root tips with extensive surface cork development. The shoot growing point of seedling not supplied boron died after five months.

Thuja plicata – Growing shoot wilts readily and the needles on young shoots become bronzed.

Pinus patula, *P. khasya* and often in *P. caribaea hondurensis* – The leading shoots became very crooked but were otherwise healthy with normal foliage and no resin flow.

Pinus elliottii – Resin flow, bud death and tip dieback symptoms. Needle malformations often occur before other external symptoms.

Boron and tree physiology

Boron is a relatively immobile nutrient within plants. Unlike many other nutrients (e.g. nitrogen and magnesium), it is not redistributed to growing points by internal cycling. Current uptake by the roots appears to determine the concentrations that are incorporated into shoots and foliage as they are formed.

Fertilization with boron (3) increases the total carbohydrate content of mycorrhizal roots. Foliar + soil fertilization yielded a 24% increase in total carbohydrates in mycorrhizal roots, whereas foliar fertilization alone decreased the total carbohydrate content. Significant increases in sugars in response to boron fertilization were observed in both ectomycorrhizal and nonmycorrhizal plants. Low boron availability limits growth of roots, and deficiency affects mycorrhizal symbiosis more than fine roots alone.

Leaves may have a light mosaic pattern of chlorosis in interveinal areas and red-brown spots interveinally and on the margins. Roots may remain short, thicken, swell into knots and break open longitudinally.

Where is boron deficiency most common?

B deficiency was most common on: soils from acid igneous rocks and fresh water sediments; acid soils from which the original content has been leached; sands low in silt, clays or micas, acid peats and mucks; and soils with free lime, including some acid soils after heavy liming (4).

Overt manifestations of B deficiency and/or reduced foliar B may be induced by addition of macronutrients (4). Liming and N fertilization (5) can cause serious B deficiency. It is likely that liming affects uptake of B, whereas N fertilization causes a dilution due to increased growth.

Environmental stresses, especially drought, often induce (7) or accentuate boron deficiency on marginally deficient sites. Although trees usually recover with resumption of normal rainfall, the incidence of multiple leaders in affected stands may be high and this reduces their economic value. Applications of boron have been observed to prevent dieback even when years of drought followed the boron application.

Soil test and plant analysis

Soil tests are most useful to determine pH and establish sufficient nutrient levels before planting, but a foliar sample (6) is needed to determine how well the tree is utilizing the soil nutrients. North Carolina State University recommends sampling at the completion of each growth flush when shoots have stopped growing and needles are fully elongated.

Studies in Australia (7) with hot-water soluble B in the 0-10 cm and 10-20 cm horizons of 0.29 and 0.19 ppm caused severe B deficiency symptoms in *P. radiata* seedlings. Soil test values are not always well correlated with plant tissue values.

Tissue concentrations					
	Leaf Boron Content, ppm B				
	Deficient	Low	Normal	High	Excess
Birch	<14		28-33		
Eastern Cottonwood	<9		68		
Eucalyptus	<35		40-70		
Holly	<20	20-25	>30		
Pinus radiata	<10			>20	
Rubber			>80		
Scots pine	<10		25-30		
Spruce	*8		25-30		

Diagnostic Criteria for Plant and Soils Ed. H.D. Chapman.
Shorrocks - Boron Deficiency and Its Cure

Boron recommendations for forestry

Christmas tree (Fraser Fir) (2) boron deficiencies have been treated with foliar application of *Solubor*[®] (1 lbs. per 100 gal.) or ground applied sprays of *Solubor*[®] (3 to 5 lbs. per acre). Do not exceed 0.25 lbs. boron per 100 gallons for foliar application to Christmas trees. Applications of 5 lbs. of boron/acre was adequate to correct B deficiency in *P. radiata* under drought conditions (7).

Soil applications (Suggested rates of application)

Crop	Granubor [®] 2 (15% B) Fertibor [®] (15% B)	Solubor [®] (20.5% B)	
	Lbs./Acre	Lbs./Acre	Gals./Acre
Birch	0.4 - 1.8	0.2 - 1.2	3 - 24
Eastern cottonwood	0.4 - 1.8	0.2 - 1.2	3 - 24
Eucalyptus	1.1 - 3.5	0.7 - 2.4	14 - 47
Pines	6.3 - 31.2	4.5 - 22.3	10.7 - 53.5
Wattle	12.5 - 37.9	8.9 - 26.8	21.4 - 64.2

References

1. Nutrient Deficiencies and Fertiliser Use in New Zealand Exotic Forests. FRI Bulletin No.97.
2. NC State University Christmas Tree Newsletter May/June 1998
3. Boron fertilization and carbohydrate relations in mycorrhizal and nonmycorrhizal shortleaf pine. Asmare Atalay et. al. 1988. *Tree Physiology* 4, 275-280.
4. Boron deficiency and excess in forest trees: A review. Earl L. Stone. *Forest Ecology and Management* 37 (1990) 49-75.
5. Reliability of Foliar Analyses of Norway Spruce Stands in a Nordic Gradient. Braekke and Salih. *Silva Fennica* 36 (2) p. 489-504 (2002)
6. Evaluating Christmas Tree Fertility. David J. Moorhead. 1996. *Georgia Christmas Tree Association Tree Talk* 10(2):14-23.
7. Boron deficiency in *Pinus radiata* D. Don and the effect of applied boron on height growth and nutrient uptake. *Plant and Soil* 79:,295-298 (1984)
8. Nutritional aspects of distorted growth in immature forest stands of southwestern coastal British Columbia. Carter and Scagel *Can J. For.*, 16, #1, 36-41 (1986)
9. Targeted Micronutrient Applications for Fraser Fir Christmas Trees. Jeffrey H. Owen. Jun 1998 *Christmas Tree Newsletter*. NC State University.
10. New Zealand Forest Research Bulletin No. 97 (Will, 1985)
11. *Forest Regeneration Manual*. Mary L. Duryea and Phillips M. Dougherty. Kluwer Academic Publishers. ISBN 0-7923-0959-6.

For more information

- Call US Borax at 1 (800) 699 9005
- Visit our website at www.borax.com/agriculture

Fertibor® Solubor® Granubor® 2